

Published Articles

Brooks S and Bailey E. (2005). Exon skipping in the KIT gene causes a Sabino spotting pattern in horses. *Mammalian Genome*, 16: 893-902.

Brooks S, Lear T, Adelson D, and Bailey E. (2007). A chromosome inversion near the KIT gene and the Tobiano spotting pattern in horses. *Cytogenetic and Genome Research*, 199: 225-230.

Brunberg E, Andersson L, Cothran G, Sandberg K, Mikko S, and Lindgren G. (2006). A missense mutation in PMEL17 is associated with the silver coat color in the horse. *BMC Genetics*, 7: 46.

Cook D, Brooks S, Bellone R, and Bailey E. (2008). Missense mutation in exon 2 of SLC36A1 responsible for champagne dilution in horses. *PLoS Genetics*, 4: e1000195.

Haase B, Brooks S, Schlumbaum A, Azor P, Bailey E, Alaeddine F, Mevissen M, Burger D, Poncet P, Rieder S, and Leeb T. (2007). Allelic heterogeneity at the equine KIT locus in dominant white (W) horses. *PLoS Genetics*, 3: e195.

Haase B, Brooks S, Tozaki T, Burger D, Poncet P, Rieder S, Hasegawa T, Penedo C, and Leeb T. (2009). Seven novel KIT mutations in horses with white coat color phenotypes. *Animal Genetics*, 40: 623-629.

Haase B, Rieder S, Tozaki T, Hasegawa T, Penedo MC, Jude R, and Leeb T. (2011). Five novel KIT mutations in horses with white coat colour phenotypes. *Animal Genetics*, 42: 337-339.

Holl H, Brooks S, and Bailey E. (2010). De novo mutation of KIT discovered as a result of a non-hereditary white coat colour pattern. *Animal Genetics*, 41: 196-198.


Mariat D, Taourit S, and Guérin G. (2003). A mutation in the MATP gene causes the cream coat colour in the horse. *Genetics, Selection, and Evolution*, 35: 119-133.

Marklund L, Johansson Moller M, Sandberg K, and Andersson L. (1996). A missense mutation in the gene for melanocyte-stimulating hormone receptor (MC1R) is associated with the chestnut coat color in horses. *Mammalian Genome*, 7: 895-899.

Metallinos D, Bowling A, and Rine J. (1998). A missense mutation in the endothelin-B receptor gene is associated with lethal white foal syndrome: an equine version of Hirschsprung Disease. *Mammalian Genome*, 9: 426-431.

Rieder S, Taourit S, Mariat D, Langlois B, and Guérin G. (2001). Mutations in the agouti (ASIP), the extension (MC1R), and the brown (TYRP1) loci and their association to coat color phenotypes in horses (*Equus caballus*). *Mammalian Genome*, 12: 450-455.

Rosengren Pielberg G, Golovko A, Sundström E, Curik I, Lennartsson J, Seltenhammer MH, Druml T, Binns M, Fitzsimmons C, Lindgren G, Sandberg K, Baumung R, Vetterlein M, Strömberg S, Grabherr M, Wade C, Lindblad-Toh K, Pontén F, Heldin CH, Sölkner J, and Andersson L. (2008) A cis-acting



regulatory mutation causes premature hair graying and susceptibility to melanoma in the horse. *Nature Genetics*, 40: 1004–1009.

Santschi E, Purdy A, Valberg S, Vrotsos P, Kaese H and Mickelson J. (1998). Endothelin receptor B polymorphism associated with lethal white foal syndrome in horses. *Mammalian Genome*, 9, 306–309.

Wagner H-J and Reissmann M. (2000). New polymorphism detected in the horse MC1R gene. *Animal Genetics*, 31: 289-290.

Yang G, Croaker D, Zhang A, Manglick P, Cartmill T, and Cass D. (1998). A dinucleotide mutation in the endothelin-B receptor gene is associated with lethal white foal syndrome (LWFS); a horse variant of Hirschsprung disease. *Human Molecular Genetics*, 7: 1047-1052.